

Steady state thermodynamics for heat conduction

Shin-ichi Sasa and Hal Tasaki

Department of Pure and Applied Sciences, University of Tokyo,
Komaba, Tokyo 153-8902, Japan, Department of Physics,
Gakushuin University, Mejiro, Toshima-ku, Tokyo 171-8588, Japan

(Dated: February 1, 2008)

Following the proposal of steady state thermodynamics (SST) by Oono and Paniconi, we develop a phenomenological theory for steady nonequilibrium states in systems with heat conduction. We find that there is essentially a unique consistent thermodynamics, and make concrete predictions, i.e, the existence of a new osmotic pressure and a shift in the coexistence temperature. These predictions allow one to test for the quantitative validity of SST by comparing them with experiments.

PACS numbers: 05.70.Ln, 44.10.-i

Construction of a statistical mechanics that apply to nonequilibrium states has been a challenging open problem in theoretical physics [1]. But so far it is not known how the desired probability measures for nonequilibrium states look like, or even whether the measures can be written in compact forms [2]. Recalling the history that the conventional thermodynamics was an essential guide when Boltzmann, Gibbs, and others constructed equilibrium statistical mechanics, it may be a good idea to start from the level of thermodynamics.

The standard theory of nonequilibrium thermodynamics [3] is based on local equilibrium hypothesis, which roughly asserts that each small part of a nonequilibrium state can be regarded as a copy of a suitable equilibrium state. But such a description seems insufficient for general nonequilibrium states. Consider, for example, a system with a steady heat flow. It is true that the quantities like the temperature and the density become essentially constant within a sufficiently small portion of the system. But no matter how small the portion is, there always exists a heat flux passing through it and hence the local state is *not* isotropic. This suggests that the local state cannot be identical to an equilibrium state (which is isotropic), but should be described rather as a *local steady state*.

Among existing attempts in nonequilibrium thermodynamics to go beyond local equilibrium treatments [4], the steady state thermodynamics (SST) proposed by Oono and Paniconi [5] seems to be most sophisticated and promising. The basic strategy of [5], in our own interpretation, is to i) look for a thermodynamics which describes a steady state as a whole, ii) clarify operational procedures for determining thermodynamic quantities, and iii) respect the general mathematical structure of thermodynamics. In the present Letter, we apply this strategy to a concrete problem of heat conduction (in a fluid), and show that theoretical consistency leads one to an essentially unique thermodynamics. We then make some concrete predictions which may be confirmed quantitatively in experiments. Extensions to other systems and discussions of related microscopic results will appear in [6].




FIG. 1: A typical system treated in SST. The left and the right walls have temperatures T and T' , respectively, and there is a steady heat flux J . We get a local steady state by restricting our attention to the thin portion within a distance L from the left wall (as denoted by the dashed line)

Local steady state: We consider a macroscopic system of a single substance in a closed cylindrical container with the cross section area A . The left and right walls of the container have very efficient heat conduction, and are kept at constant temperatures T and T' , respectively, with the aid of external heat baths. See Fig. 1. The side walls of the container are perfectly adiabatic.

If the system is kept in this setting for a sufficiently long time, it is expected to reach a unique steady state with a constant heat current without any macroscopically observable changes. (We assume that convection does not take place.) By J we denote the total energy that flow into the system from the right wall within a unit time.

We now restrict our attention to a thin portion of the system within a small distance L from the left wall as in Fig. 1. Here the length L is taken so that the temperature in the thin portion becomes essentially constant. We assume (as in local equilibrium approaches) this is realized with L which is much larger than any microscopic scales. The state in this thin portion is the *local steady state* that we study [7]. Our first crucial assumption is that the local steady state can be fully specified by four macroscopic parameters as $(T, J; V, N)$, where $V = AL$ is the volume and N is the amount of substance in the thin portion. We stress that the heat flux J need not be




FIG. 2: The porous wall in the middle of the container separates an equilibrium state and a steady nonequilibrium state. If we restrict our attention to the vicinity of the porous wall (denoted by dashed lines), we get a situation where a local steady state and an equilibrium state are in balance with each other. We use this setting to measure the chemical potential of the steady state. We will also see that there appears a force (of nonequilibrium origin) that pushes the porous wall towards the equilibrium region irrespectively to the sign of $J \neq 0$. This is an example of “flux induced osmosis.”

small.

As in the conventional thermodynamics, it is essential to consider decomposition/recombination and scaling of local steady states. In doing so, we shall always fix the cross section A and vary only the length L (within the range the system remains thin). Consider splitting the system of length L into those with lengths L_1 and L_2 (with $L = L_1 + L_2$) by a plane parallel to the left and right walls [8]. Correspondingly, we assume that the local steady state $(T, J; V, N)$ can be decomposed into two local steady states $(T, J; V_1, N_1)$ and $(T, J; V_2, N_2)$, and the two states can be recombined back into $(T, J; V, N)$. Here $V_i = AL_i$, and $N_1 + N_2 = N$. Similarly, for $\lambda > 0$, we assume that one can scale the length L to λL to get a scaled copy $(T, J; \lambda V, \lambda N)$ of the state $(T, J; V, N)$. These observations imply that the heat flux J behaves in a similar way as T , and hence should be regarded as an “intensive” variable. This identification is essential in our theory.

SST free energy: Our second essential assumption is the existence of the SST free energy $F(T, J; V, N)$, which is a thermodynamic potential describing the response of the local steady state when the extensive variables V and N are varied. We assume that there is a function $F(T, J; V, N)$ which is concave in intensive variables T, J , and convex in extensive variables V, N . It should have *additivity* $F(T, J; V, N) = F(T, J; V_1, N_1) + F(T, J; V_2, N_2)$ for any decomposition as above, *extensivity* $F(T, J; \lambda V, \lambda N) = \lambda F(T, J; V, N)$ for $\lambda > 0$, and *symmetry* $F(T, J; V, N) = F(T, -J; V, N)$ to reflect the obvious left-right symmetry of thin systems. Most importantly, we require that the derivatives of $F(T, J; V, N)$ in V and N have operational meanings exactly parallel to those in the conventional thermodynamics.

More precisely, we first require $\partial F(T, J; V, N)/\partial V = -p(T, J; V, N)$, where the pressure $p(T, J; V, N)$ is simply determined by measuring the force that the system exerts on the left (or the right) wall [9]. We then require

$\partial F(T, J; V, N)/\partial N = \mu(T, J; V, N)$, where the existence and measurability of the chemical potential $\mu(T, J; V, N)$ are assumed. A device [10] for measuring $\mu(T, J; V, N)$ is depicted in Fig. 2. Here the left and the right walls of the container are kept at constant temperatures T and T' , respectively. In the middle of the container, there is another wall made of a porous medium which is kept at a constant temperature T with the aid of an external heat bath. We assume that the substance can move across the porous wall. Suppose that the whole system has reached a steady state. Then the left half of the system has a constant temperature, and is in a normal equilibrium state (provided that the middle wall is in a very efficient contact with a bath). The right half is in a nonequilibrium steady state (which is not necessarily local) with a constant heat flux. In order to examine the balance between the two parts, we restrict our attention to a thin part of the system within a fixed distance from the porous wall as denoted by dashed lines in Fig. 2. If the distance is small enough, the part of the steady state can be regarded as local. Then we get a situation where a local steady state $(T, J; V, N)$ and an equilibrium state $(T, 0; V', N')$ are in balance with each other with respect to the exchange of the substance. Then it is natural to define $\mu(T, J; V, N)$ to be equal to $\mu(T, 0; V', N')$, where the latter can be determined within the conventional equilibrium thermodynamics. The two derivatives and the extensivity determine $F(T, J; V, N)$ without any ambiguities [11], as is obvious from the Euler equation

$$F(T, J; V, N) = -V p(T, J; V, N) + N \mu(T, J; V, N), \quad (1)$$

which is derived by using the extensivity as usual.

Let us define (extensive) SST entropy as

$$S(T, J; V, N) = -\frac{\partial F(T, J; V, N)}{\partial T}, \quad (2)$$

and a new “extensive” quantity

$$\Psi(T, J; V, N) = -\frac{\partial F(T, J; V, N)}{\partial J}, \quad (3)$$

which we shall call *nonequilibrium order parameter*. From the symmetry and the concavity of F , one finds that $\Psi(T, J; V, N) = -\Psi(T, -J; V, N) \geq 0$ for $J \geq 0$. We expect to have $\Psi(T, J; V, N) > 0$ for $J > 0$ in generic systems. Note that both S and Ψ can be measured operationally since F can be. As in the conventional thermodynamics, one can derive various identities between thermodynamic quantities.

The existence of the SST free energy with the desired properties is nothing more than an optimistic assumption. It is possible in principle that such a theoretical framework as SST simply does not exist in Nature. We therefore predict two concrete phenomena — the existence of a new osmotic pressure called flux-induced osmosis (FIO) and a shift of coexistence temperature — and present some exact relations, which enable one to test for the *quantitative* validity of SST through experiments.

Flux-induced osmosis: Let us consider the situation in Fig. 2, and examine the behavior of the pressure $p_{ss} = p(T, J; V, N)$ of the steady state. We fix the temperature T and the pressure $p_{eq} = p(T, 0; V', N')$ (by suitably varying the volume V') of the equilibrium state, and vary only the flux J . The chemical potential of the equilibrium state is thus constant, and so is the chemical potential $\mu(T, J; V, N)$ of the local steady state (by definition). We differentiate the Euler equation (1) by J with T and p_{eq} fixed. Keeping in mind that V and N may depend on J , we find

$$\frac{\partial p_{ss}(T, p_{eq}, J)}{\partial J} = \frac{\Psi(T, J; V, N)}{V}. \quad (4)$$

Noting that $p_{ss}(T, p_{eq}, 0) = p_{eq}$ and recalling the sign of Ψ , this implies $p_{ss} \geq p_{eq}$ in general. If Ψ is nonvanishing (as we expect) then one has $p_{ss} > p_{eq}$ for $J \neq 0$. Recalling that the pressures are defined from mechanical forces exerted on the walls, this implies that *the porous wall is actually pushed towards the equilibrium region irrespectively to the sign of the heat flux J* . We stress that this force, which is absent in equilibrium or local-equilibrium treatments, is of purely nonequilibrium origin [12]. This is an example of a general phenomenon that we call *flux-induced osmosis* (FIO) [6].

In the same situation, one can also derive [6] a non-trivial identity

$$\frac{\partial p_{ss}(T, p_{eq}, J)}{\partial p_{eq}} = \frac{v_{eq}}{v_{ss}}, \quad (5)$$

where $v_{eq} = V'/N'$ and $v_{ss} = V/N$. Since the identity involves only directly measurable quantities, it may be useful in quantitative tests of SST in (real and numerical) experiments.

Shift of coexistence temperature: Consider again the original setting in Fig. 1, and assume that the pressure p is kept constant. We further assume that a phase coexistence takes place in the system, i.e., the lower temperature region of the container is occupied by one phase (e.g. liquid) while the higher temperature region by another (e.g. gas). We then ask what is the temperature $T_c(p, J)$ at the boundary between the two phases. Within local equilibrium treatments, one simply concludes that $T_c(p, J)$ is the same as its equilibrium value $T_c(p, 0)$. In SST, however, we find (from an analysis similar to that for FIO) that

$$\frac{\partial T_c(p, J)}{\partial J} = -\frac{\psi_{high} - \psi_{low}}{s_{high} - s_{low}}, \quad (6)$$

where ψ_{low} and ψ_{high} are the molar Ψ of the two phases at the coexistence point, and s_{low} and s_{high} are the corresponding molar SST entropies. The identity (6) means that *in general $T_c(p, J)$ shifts from its equilibrium value*. In case $T_c(p, J) > T_c(p, 0)$ [13], one has a remarkable phenomenon of “heat flux induced freezing”, i.e., one observes a solid phase in a system with one wall having a temperature slightly higher than the melting point and the other wall having a much higher temperature.

Choice of nonequilibrium variable: Finally let us make an important remark about the choice of nonequilibrium thermodynamic variable. A local steady state may also be specified as $(T; V, N, \tau)$, where τ is the (small) temperature difference between the left and right ends of the system. Since τ is proportional to the length L in a thin uniform state, it should be regarded as an “extensive” quantity. Then, in any mathematically “healthy” thermodynamics, the corresponding free energy $\tilde{F}(T; V, N, \tau)$ (if exists) should be convex in V , N , and τ .

Following the standard argument, the convexity and the extensivity implies a variational principle

$$\begin{aligned} & \tilde{F}(T; V_1, N_1, \tau^*) + \tilde{F}(T; V_2, N_2, \tau - \tau^*) \\ & = \min_{\tau'} \{ \tilde{F}(T; V_1, N_1, \tau') + \tilde{F}(T; V_2, N_2, \tau - \tau') \}, \end{aligned} \quad (7)$$

for fixed V_1 , V_2 , N_1 , N_2 , and τ . This relation determines the temperature $T + \tau^*$ at the boundary of the two parts of the system (with volumes V_1 and V_2 , respectively) when the total temperature difference τ is fixed. In a differential form, this condition becomes $\nu(T; V_1, N_1, \tau^*) = \nu(T; V_2, N_2, \tau - \tau^*)$, where $\nu(T; V, N, \tau) = \partial \tilde{F}(T; V, N, \tau) / \partial \tau$.

On the other hand, we already know from the energy conservation law that τ^* can be determined by the condition $J(T; V_1, N_1, \tau^*) = J(T; V_2, N_2, \tau - \tau^*)$, where we expressed the heat flux J as a function of $(T; V, N, \tau)$. Since the conditions written in ν and in J must be equivalent, we conclude that there is a function f such that $J(T; V, N, \tau) = f(\nu(T; V, N, \tau))$ for any $(T; V, N, \tau)$. But, noting that J has a dimension of energy divided by time while ν is dimensionless, we find that such a universal function f simply does not exist. This observation implies that *one can never get a consistent thermodynamics by using the temperature difference τ as a nonequilibrium variable*. In fact we believe that the representation in terms of $(T, J; V, N)$, where the nonequilibrium variable J directly reflects the energy conservation law, provides an essentially unique consistent thermodynamics for a system with steady heat flux [14].

Discussions: By following the philosophy of [5], we are led to an essentially unique thermodynamics (SST) for heat conduction. The uniqueness suggests that, if a consistent thermodynamics for steady heat conduction exists at all, then it should be equivalent to what we have described here. The resulting theory led us to novel predictions which allow one to test for the quantitative validity of SST through experiments, other theoretical results [15], and numerical calculations [16].

Let us stress that SST is by no means incompatible with the long range (spatial) correlation generically found in nonequilibrium steady states [17]. The long range correlation comes from the global description based on fluctuating hydrodynamics, while local aspects are conventionally described as local steady states. The aim of SST is to provide a more sophisticated alternative of the latter, i.e., to give a unified description of nonequilibrium

modifications of the (local) equations of states. Since there is a big separation in the length scales, the existence of the long range correlation does not conflict with the validity of SST.

Perhaps the most crucial point about the potential significance of SST is whether it becomes a useful guide in the (future) construction of statistical mechanics for steady nonequilibrium states. The fact that we have arrived at an essentially unique theory is rather encouraging. We hope that, by trying to construct a statistical

theory that is consistent with the (unique) nonequilibrium thermodynamics, we are naturally led to a meaningful and correct statistical mechanics for steady nonequilibrium states.

It is a pleasure to thank Yoshi Oono for introducing us to the present subject and for various indispensable advices and suggestions. The work of S. S. was supported by grants from the Ministry of Education, Science, Sports and Culture of Japan, Nos. 12834005 and 11CE2006.

-
- [*] Electronic address: sasa@jiro.c.u-tokyo.ac.jp, hal.tasaki@gakushuin.ac.jp
- [1] See, for example, G. L. Eyink, J. L. Lebowitz, and H. Spohn, *J. Stat. Phys.* **83**, 385, (1996) and references therein.
- [2] Zubarev (*Nonequilibrium Statistical Thermodynamics*, Consultants Bureau, New York, 1974) proposed a nonequilibrium measure. But, as far as we understand, its validity has not yet been confirmed empirically. We also question whether nonequilibrium states can be described starting from local equilibrium states.
- [3] S. R. de Groot and P. Mazur, *Nonequilibrium Thermodynamics*, (Dover, New York, 1984).
- [4] See, for example, D. Jou, J. Casas-Vázquez, and G. Lebon, *Rep. Prog. Phys.* **62**, 1035 (1999) and references therein. As far as we understand, these existing attempts, although being quite suggestive and illuminating, lack theoretical rigor that we impose in our approach.
- [5] Y. Oono and M. Paniconi, *Prog. Theor. Phys. Suppl.* **130**, 29 (1998).
- [6] S. Sasa and H. Tasaki, in preparation.
- [7] We have here restricted ourselves to the simplest geometry. We plan to treat a more general class of steady states by “connecting” together local steady states.
- [8] One may regard the plane as being purely fictitious, or as a highly conducting wall kept at a suitable temperature with the aid of a heat bath.
- [9] Note that the forces exerted on the left and the right walls are exactly the same (even when the system is not “thin”). Otherwise the system gets a finite acceleration and cannot be steady.
- [10] This is not the unique way of determining μ . Another natural method is to put a system into a direct contact with a particle bath.
- [11] Our assumption here is that $F(T, J; V, N)$ with all the desired properties exists. This can be regarded as an expression of the second law for SST. Although a version of the second law is known to hold in a general Markov process (see, for example, T.M. Cover and J.A. Thomas, *Elements of Information Theory*, Wiley 1991), it is not at all clear if this second law is the one desired in SST. See also T. Hatano and S. Sasa *Phys. Rev. Lett.* **86**, 3463, (2001) and [6]. Recently Derrida, Lebowitz and Speer [See *Phys. Rev. Lett.* **89** 030601 (2002) and references therein.] studied the large deviation functionals in exclusion processes and obtained highly nontrivial and suggestive results. A large deviation functional is related to a thermodynamic free energy in equilibrium, but nothing is known for nonequilibrium. It is quite challenging to investigate possible relations between the two.
- [12] Recently Kim and Hayakawa [cond-mat/0202003] studied the Boltzmann equation and confirmed (among other things) the existence of FIO, but agreement with our predictions were only qualitative. We remark that there is a room for reexamining the determination of μ .
- [13] The concavity of the Gibbs free energy implies $s_{\text{low}} < s_{\text{high}}$ in general. We further expect to have $\psi_{\text{low}} > \psi_{\text{high}}$ in many systems.
- [14] In fact one can think about another thermodynamics in which one fixes the length L and vary only the cross section A . Then one should regard the temperature difference τ as “intensive” and its conjugate ν as “extensive.” But this theory may not be as robust as the one we propose here [6].
- [15] See [12]. We were also able to confirm the existence of SST in some stochastic models in certain limits [6]. There is also an attempt to construct a phenomenological fluctuation theory corresponding to SST (Wada and Sasa, in preparation).
- [16] Most of the “standard” stochastic models used to study nonequilibrium phenomena are designed merely to recover equilibrium distributions and fluctuations around equilibria. They may not be realistic in far-from-equilibrium situations that we are concerned with. To check the validity of SST using numerical experiments is therefore not as straightforward as one might first imagine. Nevertheless preliminary numerical analysis in the driven lattice gas is rather promising (Hayashi and Sasa, in preparation).
- [17] J.R. Dorfman, T.R. Kirkpatrick, and J.V. Sengers, *Ann. Rev. Phys. Chem.* **45**, 213 (1994) and references therein.